

The Complexity of Coverability in ν -Petri Nets

R. Lazić S. Schmitz

Department of Computer Science, U. Warwick
LSV, ENS Cachan & INRIA, U. Paris-Saclay

LICS 2016, July 6th, 2016

OUTLINE

ν -Petri nets (ν PN)

Petri nets with data management and creation

(Rosa-Velardo and de Frutos-Escrig, 2008, 2011)

coverability

- ▶ decidable by classical **backward coverability** algorithm (Abdulla et al., 2000)
- ▶ dual view using **downwards-closed** sets (Lazić and S., 2015)

complexity ν PN coverability is complete for **double Ackermann** ($\mathbb{F}_{\omega \cdot 2}$ -complete)

OUTLINE

ν -Petri nets (ν PN)

Petri nets with data management and creation

(Rosa-Velardo and de Frutos-Escrig, 2008, 2011)

coverability

- ▶ decidable by classical **backward coverability** algorithm (Abdulla et al., 2000)
- ▶ dual view using **downwards-closed** sets (Lazić and S., 2015)

complexity ν PN coverability is complete for **double Ackermann** ($F_{\omega \cdot 2}$ -complete)

OUTLINE

ν -Petri nets (ν PN)

Petri nets with data management and creation

(Rosa-Velardo and de Frutos-Escrig, 2008, 2011)


coverability

- ▶ decidable by classical **backward coverability** algorithm (Abdulla et al., 2000)
- ▶ dual view using **downwards-closed** sets (Lazić and S., 2015)

complexity ν PN coverability is complete for **double Ackermann** ($\mathbf{F}_{\omega \cdot 2}$ -complete)

v-PETRI NETS

TOKENS CARRY DATA FROM AN INFINITE COUNTABLE DOMAIN \mathbb{D}




CONFIGURATIONS IN $(\mathbb{N}^P)^\oplus$: MULTISSETS OF MARKINGS

$$\left[\begin{array}{c} \left(\begin{array}{c} 1 \\ 0 \\ 0 \end{array} \right) \\ \left(\begin{array}{c} 3 \\ 0 \\ 0 \end{array} \right) \\ \left(\begin{array}{c} 1 \\ 1 \\ 0 \end{array} \right) \end{array} \right]$$

v-PETRI NETS


TOKENS CARRY DATA FROM AN INFINITE COUNTABLE DOMAIN \mathbb{D}



CONFIGURATIONS IN $(\mathbb{N}^P)^\oplus$: MULTISSETS OF MARKINGS


$$\left[\begin{array}{c} \left(\begin{array}{c} 1 \\ 0 \\ 0 \end{array} \right) \\ \left(\begin{array}{c} 3 \\ 0 \\ 0 \end{array} \right) \\ \left(\begin{array}{c} 1 \\ 1 \\ 0 \end{array} \right) \end{array} \right] \xrightarrow{t} \left[\begin{array}{c} \left(\begin{array}{c} 1 \\ 0 \\ 0 \end{array} \right) \\ \left(\begin{array}{c} 1 \\ 1 \\ 1 \end{array} \right) \\ \left(\begin{array}{c} 0 \\ 0 \\ 0 \end{array} \right) \\ \left(\begin{array}{c} 0 \\ 0 \\ 1 \end{array} \right) \end{array} \right]$$

PETRI NETS AS ν -PETRI NETS




- ▶ a and \bar{a} are complementary **addressing** places
- ▶ c holds the actual token counts

PETRI NETS AS ν -PETRI NETS




- ▶ a and \bar{a} are complementary **addressing** places
- ▶ c holds the actual token counts

RESET PETRI NETS AS ν -PETRI NETS



- ▶ a and \bar{a} are complementary addressing places for **active** tokens
- ▶ c holds both the active and inactive tokens

RESET PETRI NETS AS ν -PETRI NETS



- ▶ a and \bar{a} are complementary addressing places for **active** tokens
- ▶ c holds both the active and inactive tokens

COVERABILITY PROBLEM

verification of safety properties “nothing bad happens”

ordering of configurations by multiset embedding

$$[\mathbf{u}_1, \dots, \mathbf{u}_n] \sqsubseteq [\mathbf{v}_1, \dots, \mathbf{v}_p]$$

iff $\exists f : \{1, \dots, n\} \rightarrow \{1, \dots, p\}$ injective ,

$\forall 1 \leq i \leq n, \mathbf{u}_i \leq \mathbf{v}_{f(i)}$

Example:


$$\left[\begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \begin{pmatrix} 3 \\ 0 \\ 0 \end{pmatrix} \right] \sqsubseteq \left[\begin{pmatrix} 10 \\ 1 \\ 0 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \begin{pmatrix} 2 \\ 0 \\ 3 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} \right]$$

input a vPN, a source configuration src , and a “bad” configuration tgt

question $\exists m, \text{tgt} \sqsubseteq m$ and $\text{src} \rightarrow^* m$?

FAST-GROWING COMPLEXITY

(S., 2016)



- ▶ Ackermann: “Ackermannian in” $x \mapsto 2x$


$$A_1(x) \stackrel{\text{def}}{=} 2x \quad A_{k+2}(x) \stackrel{\text{def}}{=} A_{k+1}^x(1) \quad A_\omega(x) \stackrel{\text{def}}{=} A_{x+1}(x)$$

- ▶ double Ackermann: “Ackermannian in” $A_\omega(x)$

$$A_{\omega+k+1}(x) \stackrel{\text{def}}{=} A_{\omega+k}^x(1) \quad A_{\omega \cdot 2}(x) \stackrel{\text{def}}{=} A_{\omega+x+1}(x)$$

FAST-GROWING COMPLEXITY

(S., 2016)



- ▶ Ackermann: “Ackermannian in” $x \mapsto 2x$


$$A_1(x) \stackrel{\text{def}}{=} 2x \quad A_{k+2}(x) \stackrel{\text{def}}{=} A_{k+1}^x(1) \quad A_\omega(x) \stackrel{\text{def}}{=} A_{x+1}(x)$$

- ▶ double Ackermann: “Ackermannian in” $A_\omega(x)$

$$A_{\omega+k+1}(x) \stackrel{\text{def}}{=} A_{\omega+k}^x(1) \quad A_{\omega.2}(x) \stackrel{\text{def}}{=} A_{\omega+x+1}(x)$$

FAST-GROWING COMPLEXITY

(S., 2016)



- ▶ Ackermann: “Ackermannian in” $x \mapsto 2x$

$$A_1(x) \stackrel{\text{def}}{=} 2x \quad A_{k+2}(x) \stackrel{\text{def}}{=} A_{k+1}^x(1) \quad A_\omega(x) \stackrel{\text{def}}{=} A_{x+1}(x)$$

- ▶ double Ackermann: “Ackermannian in” $A_\omega(x)$

$$A_{\omega+k+1}(x) \stackrel{\text{def}}{=} A_{\omega+k}^x(1) \quad A_{\omega.2}(x) \stackrel{\text{def}}{=} A_{\omega+x+1}(x)$$

MAIN RESULT

THEOREM

Coverability in ν PNs is $\mathbb{F}_{\omega,2}$ -complete.

lower bound extends Lipton's "object-oriented"
programming in Petri nets

- ▶ improves on the \mathbb{F}_{ω} lower bound of Schnoebelen (2010) for reset Petri nets
- ▶ basic block: Ackermann counters using Schnoebelen's construction
- ▶ pushed to double Ackermann: composition and iteration operations

upper bound analyses a dual view of the backward
coverability algorithm

- ▶ improves on the $\mathbb{F}_{\omega^{\omega}}$ upper bound of Rosa-Velardo (2014) for unordered data nets

MAIN RESULT

THEOREM

Coverability in ν PNs is $\mathbb{F}_{\omega,2}$ -complete.

lower bound extends Lipton's "object-oriented"
programming in Petri nets

- ▶ improves on the \mathbb{F}_{ω} lower bound of Schnoebelen (2010) for reset Petri nets
- ▶ basic block: Ackermann counters using Schnoebelen's construction
- ▶ pushed to double Ackermann: composition and iteration operations

upper bound analyses a dual view of the backward
coverability algorithm

- ▶ improves on the $\mathbb{F}_{\omega^{\omega}}$ upper bound of Rosa-Velardo (2014) for unordered data nets

v-PETRI NETS ARE WELL-STRUCTURED

(FINKEL AND SCHNOEBELEN, 2001; ABDULLA et al., 2000)

1. $((\mathbb{N}^P)^\oplus, \sqsubseteq)$ is a **well-quasi-order (wqo)**, which entails

 - finite bad sequences any sequence m_0, m_1, m_2, \dots with $\forall i < j, m_i \not\sqsubseteq m_j$, is finite
 - finite basis property any upwards-closed subset U has a finite basis B such that $U = \uparrow B$
 - ascending chain property all the ascending chains $U_0 \subsetneq U_1 \subsetneq U_2 \subsetneq \dots$ of upwards-closed subsets are finite
2. **compatibility**: if $m_1 \sqsubseteq m'_1$ and $m_1 \rightarrow m_2$, then there exists $m'_2, m_2 \sqsubseteq m'_2$ and $m'_1 \rightarrow m'_2$

v-PETRI NETS ARE WELL-STRUCTURED

(FINKEL AND SCHNOEBELEN, 2001; ABDULLA et al., 2000)

- $((\mathbb{N}^P)^\oplus, \sqsubseteq)$ is a **well-quasi-order** (wqo), which entails

 - finite bad sequences any sequence m_0, m_1, m_2, \dots with $\forall i < j, m_i \not\sqsubseteq m_j$, is finite
 - finite basis property any upwards-closed subset U has a finite basis B such that $U = \uparrow B$
 - ascending chain property all the ascending chains $U_0 \subsetneq U_1 \subsetneq U_2 \subsetneq \dots$ of upwards-closed subsets are finite
- compatibility**: if $m_1 \sqsubseteq m'_1$ and $m_1 \rightarrow m_2$, then there exists $m'_2, m_2 \sqsubseteq m'_2$ and $m'_1 \rightarrow m'_2$

"CLASSICAL" BACKWARD COVERABILITY

(ABDULLA et al., 2000)

compute $U_* \stackrel{\text{def}}{=} \bigcup_k U_k$

where

$$U_k \stackrel{\text{def}}{=} \{m' \mid \exists m \supseteq \text{tgt}, m' \rightarrow^{\leq k} m\}$$

initially $U_0 \stackrel{\text{def}}{=} \uparrow \text{tgt}$

step $U_{k+1} \stackrel{\text{def}}{=} \text{Pre}_{\exists}(U_k) \cup U_k$

where

$$\text{Pre}_{\exists}(S) \stackrel{\text{def}}{=} \{m \mid \exists s \in S, m \rightarrow s\}$$

representation of upwards-closed subsets U through their minimal elements thanks to finite basis property

termination guaranteed by ascending chain property

"CLASSICAL" BACKWARD COVERABILITY

(ABDULLA et al., 2000)

compute $U_* \stackrel{\text{def}}{=} \bigcup_k U_k$

where

$$U_k \stackrel{\text{def}}{=} \{m' \mid \exists m \supseteq \text{tgt}, m' \rightarrow^{\leq k} m\}$$

initially $U_0 \stackrel{\text{def}}{=} \uparrow \text{tgt}$

step $U_{k+1} \stackrel{\text{def}}{=} \text{Pre}_{\exists}(U_k) \cup U_k$

where

$$\text{Pre}_{\exists}(S) \stackrel{\text{def}}{=} \{m \mid \exists s \in S, m \rightarrow s\}$$

representation of upwards-closed subsets U through their minimal elements thanks to finite basis property

termination guaranteed by ascending chain property

“CLASSICAL” BACKWARD COVERABILITY

(ABDULLA et al., 2000)

compute $U_* \stackrel{\text{def}}{=} \bigcup_k U_k$

where

$$U_k \stackrel{\text{def}}{=} \{m' \mid \exists m \supseteq \text{tgt}, m' \rightarrow^{\leq k} m\}$$

initially $U_0 \stackrel{\text{def}}{=} \uparrow \text{tgt}$

step $U_{k+1} \stackrel{\text{def}}{=} \text{Pre}_{\exists}(U_k) \cup U_k$

where

$$\text{Pre}_{\exists}(S) \stackrel{\text{def}}{=} \{m \mid \exists s \in S, m \rightarrow s\}$$

representation of upwards-closed subsets U through their minimal elements thanks to finite basis property

termination guaranteed by ascending chain property

DUAL BACKWARD COVERABILITY

(LAZIĆ AND S., 2015)

compute $D_* = \bigcap_k D_k$

where

$$D_k = \{m' \mid \forall m \sqsupseteq \text{tgt}, m' \not\rightarrow^k m\}$$

initially $D_0 \stackrel{\text{def}}{=} (\mathbb{N}^P)^\otimes \setminus (\uparrow \text{tgt})$

step $D_{k+1} \stackrel{\text{def}}{=} \text{Pre}_\forall(D_k) \cap D_k$

where

$$\text{Pre}_\forall(S) \stackrel{\text{def}}{=} \{m \mid \forall s, m \rightarrow s \implies s \in S\}$$

representation of downwards-closed subsets D through
finite representations of their **ideal**
decompositions (next slide)

termination guaranteed by descending chain property

DUAL BACKWARD COVERABILITY

(LAZIĆ AND S., 2015)

compute $D_* = \bigcap_k D_k$

where

$$D_k = \{m' \mid \forall m \sqsupseteq \text{tgt}, m' \not\rightarrow^{\leq k} m\}$$

initially $D_0 \stackrel{\text{def}}{=} (\mathbb{N}^P)^\otimes \setminus (\uparrow \text{tgt})$

step $D_{k+1} \stackrel{\text{def}}{=} \text{Pre}_\forall(D_k) \cap D_k$

where

$$\text{Pre}_\forall(S) \stackrel{\text{def}}{=} \{m \mid \forall s, m \rightarrow s \implies s \in S\}$$

representation of downwards-closed subsets D through
finite representations of their **ideal**
decompositions (next slide)

termination guaranteed by descending chain property

DUAL BACKWARD COVERABILITY

(LAZIĆ AND S., 2015)

compute $D_* = \bigcap_k D_k$

where

$$D_k = \{m' \mid \forall m \sqsupseteq \text{tgt}, m' \not\rightarrow^k m\}$$

initially $D_0 \stackrel{\text{def}}{=} (\mathbb{N}^P)^\otimes \setminus (\uparrow \text{tgt})$

step $D_{k+1} \stackrel{\text{def}}{=} \text{Pre}_\forall(D_k) \cap D_k$

where

$$\text{Pre}_\forall(S) \stackrel{\text{def}}{=} \{m \mid \forall s, m \rightarrow s \implies s \in S\}$$


representation of downwards-closed subsets D through
finite representations of their **ideal**
decompositions (next slide)

termination guaranteed by descending chain property

IDEAL DECOMPOSITIONS FOR A WQO (X, \leq)

(BONNET, 1975; FINKEL AND GOUBAULT-LARRECQ, 2009)


- ▶ Downward closure
 $\downarrow S \stackrel{\text{def}}{=} \{x \in X \mid \exists s \in S. x \leq s\}$
- ▶ Ideal I
 downwards-closed, non-empty
 and directed



IDEAL DECOMPOSITIONS FOR A WQO (X, \leq)

(BONNET, 1975; FINKEL AND GOUBAULT-LARRECQ, 2009)


- ▶ Downward closure
 $\downarrow S \stackrel{\text{def}}{=} \{x \in X \mid \exists s \in S. x \leq s\}$
- ▶ Ideal I
 downwards-closed, non-empty
 and **directed**:
 $\forall x, y \in I, \exists z. x \leq z \text{ and } y \leq z$
- ▶ Examples
 - ▶ $\downarrow x \in \text{Idl}(X)$ for any x in X
 - ▶ $\mathbb{N} \in \text{Idl}(\mathbb{N})$
 - ▶ $D^\circ \in \text{Idl}(X^\circ)$ for any $D \subseteq X$
 downwards-closed



IDEAL DECOMPOSITIONS FOR A WQO (X, \leq)

(BONNET, 1975; FINKEL AND GOUBAULT-LARRECQ, 2009)


- ▶ Downward closure
 $\downarrow S \stackrel{\text{def}}{=} \{x \in X \mid \exists s \in S. x \leq s\}$
- ▶ Ideal I
 downwards-closed, non-empty
 and **directed**:
 $\forall x, y \in I, \exists z. x \leq z \text{ and } y \leq z$
- ▶ Examples
 - ▶ $\downarrow x \in \text{Idl}(X)$ for any x in X
 - ▶ $\mathbb{N} \in \text{Idl}(\mathbb{N})$
 - ▶ $D^\circ \in \text{Idl}(X^\circ)$ for any $D \subseteq X$
 downwards-closed



IDEAL DECOMPOSITIONS FOR A WQO (X, \leq)

(BONNET, 1975; FINKEL AND GOUBAULT-LARRECQ, 2009)

- ▶ Downward closure
 $\downarrow S \stackrel{\text{def}}{=} \{x \in X \mid \exists s \in S. x \leq s\}$
- ▶ Ideal I
 downwards-closed, non-empty
 and directed:
 $\forall x, y \in I, \exists z. x \leq z \text{ and } y \leq z$
- ▶ Examples
 - ▶ $\downarrow x \in \text{Idl}(X)$ for any x in X
 - ▶ $\mathbb{N} \in \text{Idl}(\mathbb{N})$
 - ▶ $D^\oplus \in \text{Idl}(X^\oplus)$ for any $D \subseteq X$
 downwards-closed




IDEAL DECOMPOSITIONS FOR A WQO (X, \leq)

(BONNET, 1975; FINKEL AND GOUBAULT-LARRECQ, 2009)

- Downward closure
 $\downarrow S \stackrel{\text{def}}{=} \{x \in X \mid \exists s \in S. x \leq s\}$

- Ideal I
 downwards-closed, non-empty
 and directed:
 $\forall x, y \in I, \exists z. x \leq z \text{ and } y \leq z$


- Examples
 - $\downarrow x \in \text{Idl}(X)$ for any x in X
 - $\mathbf{N} \in \text{Idl}(\mathbf{N})$
 - $D^\oplus \in \text{Idl}(X^\oplus)$ for any $D \subseteq X$
 downwards-closed



IDEAL DECOMPOSITIONS FOR A WQO (X, \leq)

(BONNET, 1975; FINKEL AND GOUBAULT-LARRECQ, 2009)


- ▶ Downward closure
 $\downarrow S \stackrel{\text{def}}{=} \{x \in X \mid \exists s \in S. x \leq s\}$
- ▶ Ideal I
downwards-closed, non-empty
and directed:
 $\forall x, y \in I, \exists z. x \leq z \text{ and } y \leq z$
- ▶ Examples
 - ▶ $\downarrow x \in \text{Idl}(X)$ for any x in X
 - ▶ $\mathbb{N} \in \text{Idl}(\mathbb{N})$
 - ▶ $D^\oplus \in \text{Idl}(X^\oplus)$ for any $D \subseteq X$
downwards-closed



IDEAL DECOMPOSITIONS FOR A WQO (X, \leq)

(BONNET, 1975; FINKEL AND GOUBAULT-LARRECQ, 2009)


- ▶ Downward closure
 $\downarrow S \stackrel{\text{def}}{=} \{x \in X \mid \exists s \in S. x \leq s\}$
- ▶ Ideal I
downwards-closed, non-empty
and directed
- ▶ Canonical Decompositions
if $D \subseteq X$ is downwards-closed,
then $D = I_1 \cup \dots \cup I_n$



IDEAL DECOMPOSITIONS FOR A WQO (X, \leq)

(BONNET, 1975; FINKEL AND GOUBAULT-LARRECQ, 2009)


- ▶ Downward closure
 $\downarrow S \stackrel{\text{def}}{=} \{x \in X \mid \exists s \in S. x \leq s\}$
- ▶ Ideal I
downwards-closed, non-empty
and directed
- ▶ Canonical Decompositions
if $D \subseteq X$ is downwards-closed,
then $D = I_1 \cup \dots \cup I_n$



IDEAL DECOMPOSITIONS FOR A WQO (X, \leq)

(BONNET, 1975; FINKEL AND GOUBAULT-LARRECQ, 2009)


- ▶ Downward closure
 $\downarrow S \stackrel{\text{def}}{=} \{x \in X \mid \exists s \in S. x \leq s\}$
- ▶ Ideal I
 downwards-closed, non-empty
 and directed
- ▶ Canonical Decompositions
 if $D \subseteq X$ is downwards-closed,
 then $D = I_1 \cup \dots \cup I_n$



IDEAL DECOMPOSITIONS FOR A WQO (X, \leq)

(BONNET, 1975; FINKEL AND GOUBAULT-LARRECQ, 2009)


- ▶ Downward closure
 $\downarrow S \stackrel{\text{def}}{=} \{x \in X \mid \exists s \in S. x \leq s\}$
- ▶ Ideal I
downwards-closed, non-empty
and directed
- ▶ Canonical Decompositions
if $D \subseteq X$ is downwards-closed,
then $D = I_1 \cup \dots \cup I_n$



IDEAL DECOMPOSITIONS FOR A WQO (X, \leq)

(BONNET, 1975; FINKEL AND GOUBAULT-LARRECQ, 2009)

- ▶ Downward closure
 $\downarrow S \stackrel{\text{def}}{=} \{x \in X \mid \exists s \in S. x \leq s\}$
- ▶ Ideal I
downwards-closed, non-empty
and directed
- ▶ Canonical Decompositions
if $D \subseteq X$ is downwards-closed,
then $D = I_1 \cup \dots \cup I_n$



EFFECTIVE IDEAL REPRESENTATIONS

(FINKEL AND GOUBAULT-LARRECQ, 2009; GOUBAULT-LARRECQ et al., 2016)

- ▶ extended markings:

$$\text{Idl}(\mathbb{N}^P) = \{\downarrow \mathbf{u} \mid \mathbf{u} \in \mathbb{N}_\omega^P\}$$

where $\mathbb{N}_\omega^P \stackrel{\text{def}}{=} (\mathbb{N} \cup \{\omega\})^P$

- ▶ extended configurations:

$$\text{Idl}((\mathbb{N}^P)^\otimes) = \{\downarrow (B, S) \mid B \in (\mathbb{N}_\omega^P)^\otimes, S \subseteq_f \mathbb{N}_\omega^P\}$$

- ▶ where $m \sqsubseteq (B, S)$ iff $\exists m' \in S^\otimes, m \sqsubseteq B \oplus m'$
- ▶ (B, S) is reduced iff S is an antichain and $\forall u \in \text{Support}(B), \forall v \in S, u \not\leq v$

EFFECTIVE IDEAL REPRESENTATIONS

(FINKEL AND GOUBAULT-LARRECQ, 2009; GOUBAULT-LARRECQ et al., 2016)

- ▶ extended markings:

$$\text{Idl}(\mathbb{N}^P) = \{\downarrow \mathbf{u} \mid \mathbf{u} \in \mathbb{N}_\omega^P\}$$

where $\mathbb{N}_\omega^P \stackrel{\text{def}}{=} (\mathbb{N} \cup \{\omega\})^P$

- ▶ extended configurations:

$$\text{Idl}((\mathbb{N}^P)^\otimes) = \{\downarrow (B, S) \mid B \in (\mathbb{N}_\omega^P)^\otimes, S \subseteq_f \mathbb{N}_\omega^P\}$$

- ▶ where $m \sqsubseteq (B, S)$ iff $\exists m' \in S^\otimes, m \sqsubseteq B \oplus m'$
- ▶ (B, S) is **reduced** iff S is an antichain and $\forall \mathbf{u} \in \text{Support}(B), \forall \mathbf{v} \in S, \mathbf{u} \not\leq \mathbf{v}$

EFFECTIVE IDEAL REPRESENTATIONS

(FINKEL AND GOUBAULT-LARRECQ, 2009; GOUBAULT-LARRECQ et al., 2016)

- ▶ extended markings:

$$\text{Idl}(\mathbb{N}^P) = \{\downarrow \mathbf{u} \mid \mathbf{u} \in \mathbb{N}_\omega^P\}$$

where $\mathbb{N}_\omega^P \stackrel{\text{def}}{=} (\mathbb{N} \cup \{\omega\})^P$

- ▶ extended configurations:

$$\text{Idl}((\mathbb{N}^P)^\otimes) = \{\downarrow (B, S) \mid B \in (\mathbb{N}_\omega^P)^\otimes, S \subseteq_f \mathbb{N}_\omega^P\}$$

- ▶ where $\mathbf{m} \sqsubseteq (B, S)$ iff $\exists \mathbf{m}' \in S^\otimes, \mathbf{m} \sqsubseteq B \oplus \mathbf{m}'$
- ▶ (B, S) is **reduced** iff S is an antichain and $\forall \mathbf{u} \in \text{Support}(B), \forall \mathbf{v} \in S, \mathbf{u} \not\leq \mathbf{v}$

EFFECTIVE IDEAL REPRESENTATIONS

(FINKEL AND GOUBAULT-LARRECQ, 2009; GOUBAULT-LARRECQ et al., 2016)

- ▶ extended markings:

$$\text{Idl}(\mathbb{N}^P) = \{\downarrow \mathbf{u} \mid \mathbf{u} \in \mathbb{N}_\omega^P\}$$



where $\mathbb{N}_\omega^P \stackrel{\text{def}}{=} (\mathbb{N} \cup \{\omega\})^P$

- ▶ extended configurations:

$$\text{Idl}((\mathbb{N}^P)^\otimes) = \{\downarrow (B, S) \mid B \in (\mathbb{N}_\omega^P)^\otimes, S \subseteq_f \mathbb{N}_\omega^P\}$$



- ▶ where $m \sqsubseteq (B, S)$ iff $\exists m' \in S^\otimes, m \sqsubseteq B \oplus m'$
- ▶ (B, S) is **reduced** iff S is an antichain and $\forall \mathbf{u} \in \text{Support}(B), \forall \mathbf{v} \in S, \mathbf{u} \not\leq \mathbf{v}$

DUAL BACKWARD COVERABILITY: EXAMPLE





$$D_0 = \downarrow(\omega, 4)$$

DUAL BACKWARD COVERABILITY: EXAMPLE





$$D_1 = \downarrow(1, 4) \cup \downarrow(\omega, 3)$$

DUAL BACKWARD COVERABILITY: EXAMPLE





$$D_2 = \downarrow(1, 4) \cup \downarrow(3, 3) \cup \downarrow(\omega, 2)$$

DUAL BACKWARD COVERABILITY: EXAMPLE





$$D_3 = \downarrow(1, 4) \cup \downarrow(3, 3) \cup \downarrow(5, 2) \cup \downarrow(\omega, 1)$$

DUAL BACKWARD COVERABILITY: EXAMPLE



$$D_4 = \downarrow(1,4) \cup \downarrow(3,3) \cup \downarrow(5,2) \cup \downarrow(7,1) \cup \downarrow(\omega,0)$$

DUAL BACKWARD COVERABILITY: EXAMPLE



$$D_5 = \downarrow(1,4) \cup \downarrow(3,3) \cup \downarrow(5,2) \cup \downarrow(7,1) \cup \downarrow(9,0) = D_*$$

CONTROLLED SEQUENCES

- ▶ consider a **norm** $\|\cdot\| : X \rightarrow \mathbb{N}$ with
 $\forall n, X_{\leq n} \stackrel{\text{def}}{=} \{x \in X \mid \|x\| \leq n\}$ finite:

$$\|u\| \stackrel{\text{def}}{=} \max_{p \in P \mid u(p) < \omega} u(p) \quad \text{for } u \in \mathbb{N}_{\omega}^P$$

$$\|B, S\| \stackrel{\text{def}}{=} \max_{u \in \text{Support}(B), v \in S} (\|B\|, \|u\|, \|v\|) \quad \text{for } \downarrow(B, S) \in \text{Idl}((\mathbb{N}^P)^{\otimes})$$

$$\|D\| \stackrel{\text{def}}{=} \max_{1 \leq i \leq n} \|B_i, S_i\| \quad \text{for } D = \downarrow(B_1, S_1) \cup \dots \cup \downarrow(B_n, S_n)$$

- ▶ consider a **control function** $g : \mathbb{N} \rightarrow \mathbb{N}$ strictly monotone and an **initial norm** $n \in \mathbb{N}$
- ▶ a sequence x_0, x_1, \dots of elements of X is
 (g, n) -controlled if $\forall i, \|x_i\| \leq g^i(n)$

strongly (g, n) -controlled if $\|x_0\| \leq n$ and
 $\forall i, \|x_{i+1}\| \leq g(\|x_i\|)$

CONTROLLED SEQUENCES

- ▶ consider a **norm** $\|\cdot\| : X \rightarrow \mathbb{N}$ with
 $\forall n, X_{\leq n} \stackrel{\text{def}}{=} \{x \in X \mid \|x\| \leq n\}$ finite:

$$\|\mathbf{u}\| \stackrel{\text{def}}{=} \max_{p \in P \mid \mathbf{u}(p) < \omega} \mathbf{u}(p) \quad \text{for } \mathbf{u} \in \mathbb{N}_{\omega}^P$$

$$\|\mathbf{B}, \mathbf{S}\| \stackrel{\text{def}}{=} \max_{\mathbf{u} \in \text{Support}(\mathbf{B}), \mathbf{v} \in \mathbf{S}} (\|\mathbf{B}\|, \|\mathbf{u}\|, \|\mathbf{v}\|) \quad \text{for } \downarrow(\mathbf{B}, \mathbf{S}) \in \text{Idl}((\mathbb{N}^P)^{\otimes})$$

$$\|\mathbf{D}\| \stackrel{\text{def}}{=} \max_{1 \leq i \leq n} \|\mathbf{B}_i, \mathbf{S}_i\| \quad \text{for } \mathbf{D} = \downarrow(\mathbf{B}_1, \mathbf{S}_1) \cup \dots \cup \downarrow(\mathbf{B}_n, \mathbf{S}_n)$$

- ▶ consider a **control function** $g : \mathbb{N} \rightarrow \mathbb{N}$ strictly monotone and an **initial norm** $n \in \mathbb{N}$
- ▶ a sequence x_0, x_1, \dots of elements of X is
 (g, n) -controlled if $\forall i, \|x_i\| \leq g^i(n)$

strongly (g, n) -controlled if $\|x_0\| \leq n$ and
 $\forall i, \|x_{i+1}\| \leq g(\|x_i\|)$

CONTROLLED SEQUENCES

- ▶ consider a **norm** $\|\cdot\| : X \rightarrow \mathbb{N}$ with
 $\forall n, X_{\leq n} \stackrel{\text{def}}{=} \{x \in X \mid \|x\| \leq n\}$ finite:

$$\|\mathbf{u}\| \stackrel{\text{def}}{=} \max_{p \in P \mid \mathbf{u}(p) < \omega} \mathbf{u}(p) \quad \text{for } \mathbf{u} \in \mathbb{N}_{\omega}^P$$

$$\|\mathbf{B}, \mathbf{S}\| \stackrel{\text{def}}{=} \max_{\mathbf{u} \in \text{Support}(\mathbf{B}), \mathbf{v} \in \mathbf{S}} (\|\mathbf{B}\|, \|\mathbf{u}\|, \|\mathbf{v}\|) \quad \text{for } \downarrow(\mathbf{B}, \mathbf{S}) \in \text{Idl}((\mathbb{N}^P)^{\otimes})$$

$$\|\mathbf{D}\| \stackrel{\text{def}}{=} \max_{1 \leq i \leq n} \|\mathbf{B}_i, \mathbf{S}_i\| \quad \text{for } \mathbf{D} = \downarrow(\mathbf{B}_1, \mathbf{S}_1) \cup \dots \cup \downarrow(\mathbf{B}_n, \mathbf{S}_n)$$

- ▶ consider a **control function** $g : \mathbb{N} \rightarrow \mathbb{N}$ strictly monotone and an **initial norm** $n \in \mathbb{N}$
- ▶ a sequence x_0, x_1, \dots of elements of X is
 (g, n) -controlled if $\forall i, \|x_i\| \leq g^i(n)$

strongly (g, n) -controlled if $\|x_0\| \leq n$ and
 $\forall i, \|x_{i+1}\| \leq g(\|x_i\|)$

LENGTH FUNCTION THEOREMS (1/3)

(FIGUEIRA et al., 2011; S. AND SCHNOEBELEN, 2012)

**FACT (LENGTH FUNCTION THEOREM FOR BAD SEQUENCES
IN \mathbb{N}_{ω}^P)**

Let $n > 0$. Any (g, n) -controlled bad sequence $e_0, e_1, \dots, e_{\ell}$ of extended markings in $(\mathbb{N}_{\omega}^P, \leq)$ has length at most “Ackermannian in” $g(\max(n, |P|))$.

LENGTH FUNCTION THEOREMS (2/3)

(LAZIĆ AND S., 2015)

- ▶ consider a descending chain $D_0 \supseteq D_1 \supseteq \dots \supseteq D_\ell$
- ▶ extract at each step $0 \leq k < \ell$ a **proper ideal** I_k from the canonical decomposition of D_k , s.t. $I_k \not\subseteq D_{k+1}$
- ▶ **bad sequence** of proper ideals $I_0, I_1, \dots, I_{\ell-1}$
- ▶ in particular, for descending chains $\downarrow S_0 \supseteq \downarrow S_1 \supseteq \dots \supseteq \downarrow S_\ell$ of antichains

COROLLARY (LENGTH FUNCTION THEOREM FOR HOARE-DESCENDING CHAINS OVER \mathbb{N}_ω^P)

Let $n > 0$. Any (g, n) -controlled descending chain $\downarrow S_0 \supseteq \downarrow S_1 \supseteq \dots \supseteq \downarrow S_\ell$ of antichains of $(\mathbb{N}_\omega^P, \leq)$ has length at most “Ackermannian in” $g(\max(n, |P|))$.

LENGTH FUNCTION THEOREMS (2/3)

(LAZIĆ AND S., 2015)

- ▶ consider a descending chain $D_0 \supsetneq D_1 \supsetneq \dots \supsetneq D_\ell$
- ▶ extract at each step $0 \leq k < \ell$ a **proper ideal** I_k from the canonical decomposition of D_k , s.t. $I_k \not\subseteq D_{k+1}$
- ▶ **bad sequence** of proper ideals $I_0, I_1, \dots, I_{\ell-1}$
- ▶ in particular, for descending chains $\downarrow S_0 \supsetneq \downarrow S_1 \supsetneq \dots \supsetneq \downarrow S_\ell$ of antichains

COROLLARY (LENGTH FUNCTION THEOREM FOR HOARE-DESCENDING CHAINS OVER \mathbb{N}_ω^P)

Let $n > 0$. Any (g, n) -controlled descending chain $\downarrow S_0 \supsetneq \downarrow S_1 \supsetneq \dots \supsetneq \downarrow S_\ell$ of antichains of $(\mathbb{N}_\omega^P, \leq)$ has length at most “Ackermannian in” $g(\max(n, |P|))$.

LENGTH FUNCTION THEOREMS (2/3)

(LAZIĆ AND S., 2015)

- ▶ consider a descending chain $D_0 \supsetneq D_1 \supsetneq \dots \supsetneq D_\ell$
- ▶ extract at each step $0 \leq k < \ell$ a **proper ideal** I_k from the canonical decomposition of D_k , s.t. $I_k \not\subseteq D_{k+1}$
- ▶ **bad sequence** of proper ideals $I_0, I_1, \dots, I_{\ell-1}$
- ▶ in particular, for descending chains $\downarrow S_0 \supsetneq \downarrow S_1 \supsetneq \dots \supsetneq \downarrow S_\ell$ of antichains

COROLLARY (LENGTH FUNCTION THEOREM FOR HOARE-DESCENDING CHAINS OVER \mathbb{N}_ω^P)

Let $n > 0$. Any (g, n) -controlled descending chain $\downarrow S_0 \supsetneq \downarrow S_1 \supsetneq \dots \supsetneq \downarrow S_\ell$ of antichains of $(\mathbb{N}_\omega^P, \leq)$ has length at most “Ackermannian in” $g(\max(n, |P|))$.

LENGTH FUNCTION THEOREMS (3/3)

- ▶ a descending chain $D_0 \supsetneq D_1 \supsetneq \dots \supsetneq D_\ell$ over $(\mathbb{N}^P)^\otimes$ is **star-monotone** if $\forall 0 \leq k < \ell - 1, \forall I_{k+1} = \downarrow(B_{k+1}, S_{k+1})$ proper ideal from the canonical decomposition of D_{k+1} , $\exists I_k = \downarrow(B_k, S_k)$ proper ideal from the canonical decomposition of D_k s.t. $\downarrow S_{k+1} \subseteq \downarrow S_k$

THEOREM (LENGTH FUNCTION THEOREM FOR STAR-MONOTONE DESCENDING CHAINS OVER $(\mathbb{N}_\omega^P)^\otimes$)

Let $n > 0$. Any strongly (g, n) -controlled star-monotone descending chain $D_0 \supsetneq D_1 \supsetneq \dots \supsetneq D_\ell$ of configurations in $(\mathbb{N}_\omega^P)^\otimes$ has length at most “double Ackermannian in” $g(\max(n, |P|))$.

WRAPPING UP

LEMMA (STRONG CONTROL FOR ν PNs)

The descending chain computed by the backward algorithm for a ν PN N and target tgt is strongly (g, n) -controlled for $g(x) \stackrel{\text{def}}{=} x + |N|$ and $n \stackrel{\text{def}}{=} \|\text{tgt}\|$.

LEMMA (ν PN DESCENDING CHAINS ARE STAR-MONOTONE)

The descending chains computed by the backward coverability algorithm for ν PNs are star-monotone.

THEOREM (UPPER BOUND)

The coverability problem for ν PNs is in $F_{\omega \cdot 2}$.

CONCLUDING REMARKS

- ▶ first “natural” decision problem complete for $\mathbf{F}_{\omega \cdot 2}$
- ▶ ideals and downwards-closed sets as **algorithmic** tools
 - ▶ here, backward analysis (Lazić and S., 2015)
 - ▶ forward analysis (Finkel and Goubault-Larrecq, 2009, 2012)
 - ▶ reachability in Petri nets (Leroux and S., 2015)
 - ▶ formal languages (Zetsche, 2015; Hague et al., 2016)
 - ▶ invariant inference (Padon et al., 2016)
 - ▶ piecewise testable separability (Goubault-Larrecq and S., 2016)

CONCLUDING REMARKS


- ▶ first “natural” decision problem complete for $\mathbf{F}_{\omega \cdot 2}$
- ▶ ideals and downwards-closed sets as **algorithmic** tools
 - ▶ here, backward analysis (Lazić and S., 2015)
 - ▶ forward analysis (Finkel and Goubault-Larrecq, 2009, 2012)
 - ▶ reachability in Petri nets (Leroux and S., 2015)
 - ▶ formal languages (Zetsche, 2015; Hague et al., 2016)
 - ▶ invariant inference (Padon et al., 2016)
 - ▶ piecewise testable separability (Goubault-Larrecq and S., 2016)

REFERENCES


- Abdulla, P.A., Čerāns, K., Jonsson, B., and Tsay, Y.K., 2000. Algorithmic analysis of programs with well quasi-ordered domains. *Inform. and Comput.*, 160(1–2):109–127. doi:10.1006/inco.1999.2843.
- Bonnet, R., 1975. On the cardinality of the set of initial intervals of a partially ordered set. In *Infinite and finite sets: to Paul Erdős on his 60th birthday*, Vol. 1, Coll. Math. Soc. János Bolyai, pages 189–198. North-Holland.
- Figueira, D., Figueira, S., Schmitz, S., and Schnoebelen, Ph., 2011. Ackermannian and primitive-recursive bounds with Dickson’s Lemma. In *Proc. LICS 2011*, pages 269–278. IEEE Press. doi:10.1109/LICS.2011.39.
- Finkel, A. and Schnoebelen, Ph., 2001. Well-structured transition systems everywhere! *Theor. Comput. Sci.*, 256(1–2):63–92. doi:10.1016/S0304-3975(00)00102-X.
- Finkel, A., McKenzie, P., and Picaronny, C., 2004. A well-structured framework for analysing Petri net extensions. *Inform. and Comput.*, 195(1–2):1–29. doi:10.1016/j.ic.2004.01.005.
- Finkel, A. and Goubault-Larrecq, J., 2009. Forward analysis for WSTS, part I: Completions. In *Proc. STACS 2009*, volume 3 of *Leibniz Int. Proc. Inf.*, pages 433–444. LZI. doi:10.4230/LIPIcs.STACS.2009.1844.
- Finkel, A. and Goubault-Larrecq, J., 2012. Forward analysis for WSTS, part II: Complete WSTS. *Logic. Meth. in Comput. Sci.*, 8(3:28):1–35. doi:10.2168/LMCS-8(3:28)2012.
- Goubault-Larrecq, J. and Schmitz, S., 2016. Deciding piecewise testable separability for regular tree languages. Preprint. hal.inria.fr:hal-01276119.
- Goubault-Larrecq, J., Karandikar, P., Narayan Kumar, K., and Schnoebelen, Ph., 2016. The ideal approach to computing closed subsets in well-quasi-orderings. In preparation. See also an earlier version in: J. Goubault-Larrecq. On a generalization of a result by Valk and Jantzen. Research Report LSV-09-09, LSV, ENS Cachan, 2009. URL http://www.lsv.fr/Publis/RAPPORTS_LSV/PDF/rr-lsv-2009-09.pdf.
- Haddad, S., Schmitz, S., and Schnoebelen, Ph., 2012. The ordinal recursive complexity of timed-arc Petri nets, data nets, and other enriched nets. In *Proc. LICS 2012*, pages 355–364. IEEE Press. doi:10.1109/LICS.2012.46.
- Hague, M., Kochems, J., and Ong, C.H.L., 2016. Unboundedness and downward closures of higher-order pushdown automata. In *POPL 2016*, pages 151–163. ACM. doi:10.1145/2837614.2837627.
- Lazić, R., Newcomb, T., Ouaknine, J., Roscoe, A., and Worrell, J., 2008. Nets with tokens which carry data. *Fund. Inform.*, 88(3):251–274.
- Lazić, R. and Schmitz, S., 2015. The ideal view on Rackoff’s coverability technique. In *Proc. RP 2015*, volume 9328 of *Lect. Notes in Comput. Sci.*, pages 1–13. Springer. doi:10.1007/978-3-319-24537-9_8.
- Leroux, J. and Schmitz, S., 2015. Demystifying reachability in vector addition systems. In *LICS 2015*, pages 56–67. IEEE Press. doi:10.1109/LICS.2015.16.

- Lipton, R., 1976. The reachability problem requires exponential space. Technical Report 62, Yale University.
- Padon, O., Immerman, N., Shoham, S., Karbyshev, A., and Sagiv, M., 2016. Decidability of inferring inductive invariants. In *POPL 2016*, pages 217–231. ACM. doi:10.1145/2837614.2837640.
- Rackoff, C., 1978. The covering and boundedness problems for vector addition systems. *Theor. Comput. Sci.*, 6(2): 223–231. doi:10.1016/0304-3975(78)90036-1.
- Rosa-Velardo, F. and de Frutos-Escrig, D., 2008. Name creation vs. replication in Petri net systems. *Fund. Inform.*, 88 (3):329–356.
- Rosa-Velardo, F. and de Frutos-Escrig, D., 2011. Decidability and complexity of Petri nets with unordered data. *Theor. Comput. Sci.*, 412(34):4439–4451. doi:10.1016/j.tcs.2011.05.007.
- Rosa-Velardo, F. and Martos-Salgado, M., 2012. Multiset rewriting for the verification of depth-bounded processes with name binding. *Inform. and Comput.*, 215:68–87. doi:10.1016/j.ic.2012.03.004.
- Rosa-Velardo, F., 2014. Ordinal recursive complexity of unordered data nets. Technical Report TR-4-14, Departamento de Sistemas Informáticos y Computación, Universidad Complutense de Madrid. <http://antares.sip.ucm.es/frosa/docs/complexityUDN.pdf>.
- Schmitz, S. and Schnoebelen, Ph., 2012. Algorithmic aspects of WQO theory. Lecture notes. <http://cel.archives-ouvertes.fr/cel-00727025>.
- Schmitz, S., 2016. Complexity hierarchies beyond Elementary. *ACM Trans. Comput. Theory*. <http://arxiv.org/abs/1312.5686>. To appear.
- Schnoebelen, Ph., 2010. Revisiting Ackermann-hardness for lossy counter machines and reset Petri nets. In *Proc. MFCS 2010*, volume 6281 of *Lect. Notes in Comput. Sci.*, pages 616–628. Springer. doi:10.1007/978-3-642-15155-2_54.
- Zetsche, G., 2015. An approach to computing downward closures. In *ICALP 2015*, volume 9135 of *Lect. Notes in Comput. Sci.*, pages 440–451. Springer. doi:10.1007/978-3-662-47666-6_35.


TAXONOMY OF PETRI NET EXTENSIONS



TAXONOMY OF PETRI NET EXTENSIONS



TAXONOMY OF PETRI NET EXTENSIONS



POLYADIC ν -PETRI NETS

(ROSA-VELARDO AND MARTOS-SALGADO, 2012)

- ▶ hold *tuples* of tokens in places
- ▶ equivalent to the full π -*calculus*
- ▶ model of *dynamic* database systems with existential positive guards
- ▶ *undecidable* coverability