

## TD 9: BDDs

**Exercise 1** (Some BDDs). Draw the BDDs for the following functions, using the order of your choice on the variables  $\{x_1, x_2, x_3\}$ :

1. the majority function  $m(x_1, x_2, x_3)$ : its value is 1 iff the majority of the input bits are 1's,
2. the hidden weighted bit function  $h(x_1, x_2, x_3)$ : its value is that of variable  $x_s$ , where  $s = \sum_{i=1}^3 x_i$  and  $x_0$  is defined as 0.

**Exercise 2** (Symmetric Functions). A *symmetric function* of  $n$  variables has the same value for all permutations of the same  $n$  tuple of arguments.

Show that a BDD for a symmetric function has at most  $\frac{n(n+1)}{2} + 1$  nodes (when omitting the 0-node).

**Exercise 3** (Counting Solutions). Write a linear time algorithm for counting the number of solutions of a boolean function  $f$  represented by a BDD, i.e. of the number of valuations  $\nu$  s.t.  $\nu \models f$ .

**Exercise 4** (An Upper Bound on the Size of BDDs). The size  $B(f)$  of a BDD for a function  $f$  is defined as the number of its nodes. Consider an arbitrary boolean function  $f$  on the ordered set  $x_1 \cdots x_n$ , and consider a variable  $x_k$ .

1. Show that we can bound the number of nodes labeled by  $\{x_1, \dots, x_k\}$  by  $2^k - 1$ .
2. How many different subfunctions on the ordered set of variables  $x_{k+1} \cdots x_n$  exist? Deduce another bound for the number of nodes labeled by  $\{x_{k+1}, \dots, x_n\}$ .
3. What global bound do you obtain for  $k = n - \log_2(n - \log_2 n)$ ?

**Exercise 5** (Finding the Optimal Order). There are in general  $n!$  different orders for the variables  $\{x_1, \dots, x_n\}$ , and building the BDD for each of these is computationally expensive. One can nevertheless design an exponential time algorithm for finding the optimal order. Indeed, an optimal ordering on a subset  $X$  of variables does not depend on the order in which  $X' = \{x_1, \dots, x_n\} \setminus X$  has been accessed.

1. Fix a boolean function  $f$  over variables  $\{x_1, \dots, x_n\}$ . We assume that  $f$  is provided as a BDD  $B$  for the ordering  $x_1, x_2, \dots, x_n$ .

Given a subset  $X$  of  $\{x_1, \dots, x_n\}$  and a variable  $x$  in  $X$ , how many nodes labeled by  $x$  does any BDD  $B'$  for  $f$  has if it first treats  $X' = \{x_1, \dots, x_n\} \setminus X$ , then  $x$ , and last  $X \setminus \{x\}$ ? How can you compute this number on the provided BDD  $B$  for  $f$ ?

2. Reduce the optimal order problem to the search of a path of minimal weight in a weighted graph with subsets of  $\{x_1, \dots, x_n\}$  as vertices.